


ICL SURGERY DROP INSTRUCTIONS

The eye will take approximately 1 month to heal after your surgery. During this time, your vision may be blurry and you may experience temporary side effects such as: a scratching and burning sensation, excessive tearing, sensitivity to light, redness, swollen eyelids, and a foreign body sensation. You should not have significant pain and if you do, please contact your eye doctor.

MEDICATION INSTRUCTIONS

RIGHT EYE/LEFT EYE

MEDICATION	PURPOSE	DOSAGE
<p>PRED – MOXI – NEPA:</p> 	<p>Reduces Pain & Inflammation and Prevents Infection</p>	<p><u>PRED – MOXI - NEPA:</u></p> <p><u>PRIOR TO SURGERY:</u> 1 DROP <u>4</u> TIMES A DAY FOR 2 DAYS -NO DROPS DAY OF SURGERY</p> <p><u>AFTER SURGERY:</u> Week 1: 1 DROP <u>4</u> TIMES A DAY Week 2: 1 DROP <u>3</u> TIMES A DAY Week 3: 1 DROP <u>2</u> TIMES A DAY Week 4: 1 DROP <u>1</u> TIME A DAY FOR 1 WEEK THEN STOP</p>
<p>Eye Shield</p>	<p>Protects eye while asleep</p>	<p>Tape over eye while sleeping for 1 week after surgery</p>
<p>Artificial Tears</p>	<p>Lubricates the eye</p>	<p>1 Drop as Needed</p>

- If using artificial tears or any other drops, please wait five minutes after applying the Pred-Gati-Brom to avoid washing out the medication.
- Avoid heavy lifting, bending, and strenuous activities until next visit.