

CATARACT POST-OPERATIVE DROP INSTRUCTIONS FOR DR. GABIANELLI, DR. FEINBERG, DR. SAWHNEY, Dr. SCHOENBERG

The eye will take approximately 1 month to heal after your surgery. During this time, your vision may be blurry and you may experience temporary side effects such as: a scratching and burning sensation, excessive tearing, sensitivity to light, redness, swollen eyelids, and a foreign body sensation. You should not have significant pain and if you do, please contact your eye doctor.

MEDICATION INSTRUCTIONS

RIGHT EYE/LEFT EYE

MEDICATION- (circle med)	PURPOSE	DOSAGE
	<p style="text-align: center;">Reduces Pain & Inflammation</p>	<p><u>PROLENSA</u> or <u>ILEVRO</u>: 1 DROP <u>1</u> A DAY FOR <u>3</u> WEEKS AFTER SURGERY THEN STOP</p>
	<p style="text-align: center;">Prevents Infection</p>	<p><u>VIGAMOX</u> or <u>BESIVANCE</u>: 1 DROP <u>4</u> TIMES A DAY FOR <u>7</u> DAYS AFTER SURGERY THEN STOP</p>
	<p style="text-align: center;">Reduces Inflammation</p>	<p><u>DUREZOL</u>: 1 DROP <u>3</u> TIMES A DAY FOR 1 WEEK, THEN 1 DROP <u>2</u> TIMES A DAY FOR 2 WEEKS, THEN 1 DROP <u>1</u> TIME A DAY FOR 1 WEEK, THEN STOP</p> <p>or</p> <p><u>PRED FORTE</u>: 1 DROP <u>4</u> TIMES A DAY FOR 1 WEEK, THEN 1 DROP <u>3</u> TIMES A DAY FOR 1 WEEK, THEN 1 DROP <u>2</u> TIMES A DAY FOR 1 WEEK, THEN 1 DROP <u>1</u> TIME A DAY FOR 1 WEEK, THEN STOP</p>
<p style="text-align: center;">Eye Shield</p>	<p style="text-align: center;">Protects eye while asleep</p>	<p style="text-align: center;">TAPE OVER EYE AT BEDTIME FOR 1 WEEK AFTER SURGERY</p>
<p style="text-align: center;">Artificial Tears</p>	<p style="text-align: center;">Lubricates the eye</p>	<p style="text-align: center;">1 DROP AS NEEDED</p>

* TO AVOID WASHING OUT THE PREVIOUSLY APPLIED DROPS, IT IS BEST TO SEPARATE DROPS BY ABOUT 5 MINUTES IN BETWEEN. AVOID HEAVY LIFTING, BENDING, AND STRENOUS ACTIVITIES UNTIL NEXT VISIT.